

UDK:-616.441:616.314.2-092

CLINICAL AND PATHOGENETIC ASPECTS OF DISEASES OF HARD TISSUES, TEETH, AND PERIODONT IN PATIENTS WITH THYROID GLANDS DISORDER.

Akhmedov A.A., Sheraliyev F.I.
Samarkand State Medical University.

ANNOTATION

Thyroid diseases are increasing worldwide. Thyroid pathology ranks second among all endocrine diseases after diabetes mellitus. This is due to the influence of various exogenous factors: iodine deficiency in the environment, the effects of various stromogenic factors, the deterioration of the ecological situation, and improved diagnostics of thyroid nodules. Studying the impact of thyroid pathology on the condition of periodontal tissues is also important because one of its diseases - endemic goiter - is quite widespread, especially in regions with a deficiency of a number of natural focal factors - iodine and fluorine. This is due to the influence of exogenous factors: iodine deficiency in the environment, the action of various stromogenic factors, the deterioration of the ecological situation, and improved diagnostics of thyroid nodules.

Keywords: periodontium, periodontal diseases, dental hard tissue diseases, somatic diseases, dental nail.

АННОТАЦИЯ

Во всем мире отмечается рост числа заболеваний щитовидной железы. Тиреоидная патология занимает второе место среди всех эндокринных заболеваний после сахарного диабета. Это связано с воздействием экзогенных факторов: недостаток йода в окружающей среде, действие различных стромогенных факторов, ухудшение экологической ситуации, улучшение диагностики узловых образований щитовидной железы. При гипотиреозе выявлена высокая распространенность воспалительных процессов пародонта. Изучение влияния патологии щитовидной железы на состояние тканей пародонта важно также и в связи с тем, что одно из ее заболеваний – эндемический зоб - имеет довольно широкое распространение, особенно в регионах с недостатком ряда природно-очаговых факторов - йода и фтора. Это связано с воздействием экзогенных факторов: недостаток йода в окружающей среде, действие различных стромогенных факторов, ухудшение экологической ситуации, улучшение диагностики узловых образований щитовидной железы.

Ключевые слова: пародонт, заболевания пародонта, заболеваний твердых тканей зубов, соматические заболевания, зубной налет, профилактика, гигиена полости рта.

ANNOTATSIYA.

Butun dunyoda qalqonsimon bez kasalliklari soni ortib bormoqda. Qalqonsimon bez patologiyasi barcha endokrin kasalliklar orasida qandli diabetdan keyin ikkinchi o'rinda turadi. Bu ekzogen omillarning ta'siri bilan bog'liq: atrof-muhitda yod tanqisligi, turli xil strumogen omillarning ta'siri, ekologik vaziyatning yomonlashishi, qalqonsimon bezning tugunli hosilalari diagnostikasini yaxshilash. Qalqonsimon bez patologiyasining parodont to'qimalari holatiga ta'sirini o'rganish uning

kasalliklaridan biri - endemik bo'qoq, ayniqsa, bir qator tabiiy o'choqli omillar - yod va fluor kam bo'lgan hududlarda keng tarqalganligi bilan ham muhimdir. Bu ekzogen omillarning ta'siri bilan bog'liq: atrof-muhitda yod tanqisligi, turli xil omillarning ta'siri, ekologik vaziyatning yomonlashishi sabab bo'ladi.

Kalit so'zlar: parodont, parodont kasalliklari, tish qattiq to'qimalari kasalliklari somatik kasalliklar, tish karashi, profilaktika, og'iz bo'shlig'i gigiyenasi.

Research objectives:

To study the condition of the hard tissues of the teeth and periodontium in patients with diffuse endemic goiter of mild and moderate severity;

to determine morphostructural changes in the microelement composition in patients with diffuse endemic goiter of mild and moderate severity using the scanning microscopy method;

to analyze the biochemical indicators of bone metabolism in oral fluid by determining the content of total calcium, phosphorus, and alkaline phosphatase in oral fluid in patients with diffuse endemic goiter of mild and moderate severity;

to study the phagocytic, cellular, and humoral immune status of patients with enamel and dentin demineralization and generalized periodontitis against the background of diffuse endemic goiter of mild and moderate severity;

Improve measures for the prevention of caries and treatment of patients with generalized periodontitis against the background of diffuse endemic goiter of mild and moderate severity.

The object of the investigation: The object of the study was 115 patients with caries and chronic generalized periodontitis against the background of diffuse endemic goiter aged 25-45 years. The control group consisted of 40 patients with caries and inflammatory diseases of the periodontium, without somatic pathology and not residing in the endemic zone.

Subject of research: oral fluid, periodontal tissues for determining dental indicators.

Table 1

Results of the Green-Vermilion oral hygiene index of the studied groups

Research methods. To achieve the goal and objectives, the following methods were used: instrumental, clinical-functional, laboratory, radiological, and statistical.

To fulfill the set goals and objectives of the study, a clinical and instrumental study of the condition of periodontal tissues was conducted in 75 patients aged 25-45 years with caries and chronic generalized periodontitis against the background of diffuse endemic goiter of I, II degree at the Samarkand Regional Stomatology Polyclinic and the Samarkand Regional Endocrinology Dispensary between 2020 and 2024. The control group consisted of 40 patients with caries and inflammatory diseases of the periodontium without thyroid pathology. In the second stage, to assess the effectiveness of the treatment and preventive measures we proposed, patients with diffuse endemic goiter were divided into 2 groups: the comparison group - 30 patients who received standard traditional treatment, and the main group - 45 patients who, along with standard treatment, received the complex treatment we proposed.

The results of the work "Clinical Observations and Analysis of Own Research" analyzed the simplified oral hygiene index (OHI-S). In patients with endemic goiter, an increase in this indicator was observed compared to the control group (Table 1). A low hygiene index of 1.2 ± 0.15 in the control group aged 25-35 indicates good oral hygiene. This index was also higher in the 35-45 age group compared to the control group (1.8 ± 0.20).

Age	Control group	Main group
25-35	1,2±0,15	1,9±0,13*
35-45	1,8±0,20	2,6±0,31*
Average parameter	1,5±0,17	2,25±0,22*

Note. * - differences are significant compared to the control group; $p < 0.05$

Loe, Silness index (table. 2) reflected the presence of inflammatory phenomena manifested as gingivitis in the 25-35 age group. Severe gingivitis prevailed in the 35-45 age group. The lowest index in the control group

was 2.22 ± 0.11 , which is 0.40 lower than in patients of the main group. The degree of gingivitis was moderate and severe in both groups. The gingival index was higher in the main group than in the control group.

Table.2

Gingival GI (Loe, Silness)

Age	Control group	Main group
25-35	1,23±0,06	1,52±0,10*
35-45	2,22± 0,11	2,62±0,14*
Average parameter	1,73±0,06	2,07±0,11*

Note. * - differences are significant compared to the control group; $p < 0.05$

During cytological examination in patients' cytograms, the main

in the group, moderate cellularity of the cytograms with a predominance of damaged tissue structures of the periodontium was noted, the absence of immunocompetent cells against a background of moderate leukocyte

infiltration, a moderate amount of fungal mycelial threads were revealed, indicating a decrease in protective processes in the periodontium during the development of chronic inflammation. In patients of the control group, immunocompetent cells were found in moderate amounts in all preparations, and fungal mycelium was detected in isolated cases.




Figure 1. Sitogram of the main group of patients with moderate periodontitis. A significant number of total nuclear elements. Romanovsky-Giemsa coloring. Increase by 200.

In the study of the dynamics of blood serum cytokine levels, significant differences were established after treatment in the main group in the indicators of IL-1RA, IL-8, $\text{TNF}\alpha$, and $\text{INF}\gamma$ (Table 18). The concentration of IL-

1RA in the main group after treatment decreased significantly by 51.8% (876.5 ± 97.35) after 1 month, and by 57.5% (774.1 ± 85.42) after 6 months from the baseline level (1743 ± 162.1). The level of IL-8 significantly decreased in the main group after treatment by 76.5% (18.22 ± 3.11) after one month and by 80.6% (14.88 ± 4.17) after 6 months from the

baseline (78.5 ± 0.03). A significant decrease in the concentration of $\text{TNF}\alpha$ was detected only after 6 months in patients of the main group by 14.8% (2.13 ± 0.07) of the baseline (2.48 ± 0.06). The dynamics of $\text{INF}\gamma$ in the form of a significant 68% increase was observed after 1 and 6 months only in patients of the main group.

The concentration of immunoglobulins of all classes in oral fluid varied significantly only in the main group. The concentration of IgA in the oral fluid is indicated as an indicator of a decrease in the inflammatory process, most pronounced after 1 month of treatment by 80.3% (0.102 ± 0.02) of the initial level (0.52 ± 0.02), which is confirmed by the positive dynamics of clinical and laboratory indicators. After 6 months, the concentration of IgG was 91.2% (0.08 ± 0.02). In the main group, the concentration of IgM in oral fluid significantly decreased from 72.6% (0.005 ± 0.002) after 1 month to 90.8% (0.003 ± 0.002) after 6 months (0.022 ± 0.007), but in the control group, a tendency to decrease was observed only after 6 months.

Conclusions. Chronic generalized periodontitis in patients with diffuse endemic goiter is characterized by a prolonged latent course with characteristic clinical manifestations in the form of a combination of generalized gingival recession, pronounced loss of clinical attachment, and a potentially long-term course. In the presence of endemic goiter, a significant increase in the frequency of dental caries occurs. Analysis of dental CPI indicators showed that in 100% of cases in the main group, there were carious lesions of the hard tissues of the teeth. In the control group, the CPI index was 9.4 ± 1.7 and significantly lower than in the group with diffuse endemic goiter.

The cytological picture in periodontitis in patients with diffuse endemic goiter is characterized by moderate leukocyte infiltration and the absence of immunocompetent cells. The BMI indicators for periodontitis in patients with hypothyroidism are 1.4 times (27.5%) higher, the BMI indicators are 1.3 times (33.3%) higher than the corresponding indicators in patients with chronic generalized periodontitis without thyroid pathology ($p < 0.05$). In the groups of patients with moderate and severe endemic goiter, a decrease in calcium content (0.26 mmol/l) and an increase in magnesium (3.4 mmol/l) against a background of slight changes in phosphorus levels (0.4 mmol/l) were found in mixed saliva. A decrease in calcium concentration undoubtedly affects the mineralizing ability of saliva, which, against the background of an increase in its acidity, will adversely affect the condition of the hard tissues of the tooth. Chronic periodontitis against the background of primary hypothyroidism is characterized by disorders in immune homeostasis: marked suppression of the factors protecting the phagocytic link of the immune system in the form of a significant decrease in the indicators of AP, NST-test, and FI; insufficiency of the cellular link of immunity, expressed in a decrease in the level of CR4-lymphocytes and the immunoregulatory index; insufficiency of the humoral link in the form of a decrease in the concentration of IgM and IgA in blood serum, a decrease in the concentration of IgM, an increase in IgA and IgG in oral fluid. The cytokine status is characterized by an increase in the concentration of IL-1PA in oral fluid, an increase in the concentration of IL-8 in blood serum, and a decrease in the level of IL-4.

REFERENCES| CHOCKH | IQTIBOSLAR:

1. Asrorovna, X. N., Baxriddinovich, T. A., Bustanovna, I. N., Valijon O'g'li, D. S., & Qizi, T. K. F. (2021). Clinical Application Of Dental Photography By A Dentist. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(09), 10-13.
2. Bustanovna, I. N. (2024). Hygienic Assessment of The Condition of The Oral Mucosa After Orthopedic Treatment. *International Journal of Scientific Trends*, 3(3), 56-61.

3. Bustanovna, I. N. (2024). Complications Arising in the Oral Cavity after Polychemotherapy in Patients with Hemablastoses. *International Journal of Scientific Trends*, 3(3), 62-66.
4. Bustanovna, P. I. N. (2024). Further Research the Features of the Use of Metal-Ceramic Structures in Anomalies of Development and Position of Teeth. *International Journal of Scientific Trends*, 3(3), 67-71.
5. Ghasemi Darestani, N., Gilmanova, A. I., Al-Gazally, M. E., Zekiy, A. O., Ansari, M. J., Zabibah, R. S., ... & Akhavan-Sigari, R. (2023). Mesenchymal stem cell-released oncolytic virus: an innovative strategy for cancer treatment. *Cell Communication and Signaling*, 21(1), 43.
6. Islamova N. B., Sh N. N. STUDY OF CHANGES IN PERIODONTAL DISEASES IN POST-MENOPAUSAL WOMEN //Conferences. – 2023. – C. 15-17.
7. Islamova, N. B. (2022). CHANGES IN PERIODONTAL TISSUES IN THE POSTMENOPAUSAL PERIOD. In *Стоматология-наука и практика, перспективы развития* (pp. 240-241).
8. Pallathadka, H., Khaleel, A. Q., Zwamel, A. H., Malathi, H., Sharma, S., Rizaev, J. A., ... & Jawad, M. A. (2024). Multi-Drug Resistance and Breast Cancer Progression via Toll-Like Receptors (TLRs) Signaling. *Cell Biochemistry and Biophysics*, 1-16.
9. Fakhridin, C. H. A. K. K. A. N. O. V., Shokhrub, S. A. M. A. D. O. V., & Nilufar, I. S. L. A. M. O. V. A. (2022). ENDOKANAL PIN-KONSTRUKSIYALARNI ISHLATISHDA ASO-RATLAR VA XATOLAR TAHLILI. *JOURNAL OF BIOMEDICINE AND PRACTICE*, 7(1).
10. Rizaev, J. A., Nazarova, N. S., & Vohidov, E. R. (2024). HOMILADOR AYOLLARDA PARODONT KASALLIKLARI RIVOJLANISHINING PATOGENETIK JIHATLARI. *Журнал гуманитарных и естественных наук*, (11 [2]), 104-107.
11. RIZAEV, J., & KUBAEV, A. (2020). Preoperative mistakes in the surgical treatment of upper retro micrognathia. *International Journal of Pharmaceutical Research (09752366)*, 12(1).
12. Kizi, J. O. A., & Bustanovna, I. N. (2024). FAMILIARIZATION WITH THE HYGIENIC ASSESSMENT OF THE CONDITION OF THE ORAL MUCOSA IN ORTHOPEDIC TREATMENT. *European International Journal of Multidisciplinary Research and Management Studies*, 4(05), 89-96.
13. Ugli, A. A. A., & Bustanovna, I. N. (2024). STUDY OF THE CONDITION OF PARODONT IN PERIODONTITIS IN FETAL WOMEN. *European International Journal of Multidisciplinary Research and Management Studies*, 4(05), 149-156.